

COMPACT CLOSED BICATEGORIES

MICHAEL STAY

ABSTRACT. A compact closed bicategory is a symmetric monoidal bicategory where every object is equipped with a weak dual. The unit and counit satisfy the usual “zig-zag” identities of a compact closed category only up to natural isomorphism, and the isomorphism is subject to a coherence law.

We give several examples of compact closed bicategories, then review previous work. In particular, Day and Street defined compact closed bicategories indirectly via Gray monoids and then appealed to a coherence theorem to extend the concept to bicategories; we restate the definition directly, emphasizing the combinatorics. Finally, we prove that each of the examples are compact closed.

We prove that given a 2-category C with finite products and weak pullbacks, the bicategory of objects of C , spans, and isomorphism classes of maps of spans is compact closed. We also prove that given a cocomplete symmetric monoidal category R whose tensor product distributes over its colimits, the bicategory $\text{Mat}(R)$ of natural numbers, matrices of objects of R and matrices of morphisms of R is compact closed. As corollaries, the bicategory of spans of sets, the bicategory of relations, and certain bicategories of “resistor networks” are all compact closed. We also give a new proof that the 2-category of small categories, cocontinuous functors between the presheaves on those categories, and natural transformations is compact closed.

1. Introduction

When moving from set theory to category theory and higher to n -category theory, we typically weaken equations at one level to natural isomorphisms at the next. These isomorphisms are then subject to new coherence equations. For example, multiplication in a monoid is associative, but the tensor product in a monoidal category is, in general, only associative up to a natural isomorphism. This “associator” natural isomorphism has to satisfy some extra equations that are trivial in the case of the monoid. In a similar way, when we move from compact closed categories to compact closed bicategories, the “zig-zag” equations governing the dual get weakened to natural isomorphisms and we need to introduce some new coherence laws.

In Section 2, we will give several examples of important mathematical structures and how they arise in relation to compact closed bicategories. Following the examples, in Section 3 we give the history of the concept and related work. Next, in Section 4 we give the complete definition, which to our knowledge has not appeared elsewhere; we try to motivate each piece of the definition so that the reader could afterwards reconstruct the definition without the aid of this paper. In Section 5, we prove that a construction by

2000 Mathematics Subject Classification: 18D05.

Key words and phrases: category theory, bicategory, compact closed.


© Michael Stay, 2012. Permission to copy for private use granted.

Hoffnung is an instance of a compact closed bicategory. We prove our assertions about the examples in Section 6. Most of these are corollaries of the main theorem.


2. Examples

In order to get across some of the flavor of these different compact closed bicategories, we will describe the bicategories as well as some weak monoids and monads in them. The weak monoids and monads play no role in the rest of the paper, but we have found that comparing and contrasting them helps when trying to develop intuition about the bicategories. The monoids are variations on the notion of an associative algebra, while the monads are variations on the notion of a category. We will give proofs that each of these bicategories (with the exception of $nCob_2$) are compact closed in Section 6.


- A **span** from X to Y in a category C is an object S in C together with an ordered pair of morphisms $(f: S \rightarrow X, g: S \rightarrow Y)$.



If C is a category with pullbacks, we can compose spans:



A **map of spans** between two spans $Y \xleftarrow{g} S \xrightarrow{f} X$ and $Y \xleftarrow{g'} S' \xrightarrow{f'} X$ is a morphism $h: S \rightarrow S'$ making the following diagram commute:



Since the pullback is associative only up to a natural isomorphism, the same is true of the composition of two spans, so this construction does not give a 2-category; however, we do get a bicategory $\text{Span}(C)$ of objects of C , spans in C , and maps of spans.

If C is a category with finite products as well as pullbacks, then the bicategory $\text{Span}(C)$ is a compact closed bicategory where the tensor product is given by the product in C . A weak monoid object in $\text{Span}(C)$ is a categorification of the notion of an associative algebra. For example, one weak monoid in $\text{Span}(\text{Set})$ is equivalent to the category of polynomial functors from Set to itself; such functors can be “added” using disjoint union, “multiplied” using the cartesian product, and “scaled” by sets [23]. A monad in $\text{Span}(C)$ is a category internal to C [10].

- Sets, relations, and implications form the compact closed bicategory Rel , where the tensor product is given by the product in Set . Let R be the ring of truth values; then a weak monoid object M in Rel is a quantale on the powerset of M [49], while a monad in Rel is a preorder.
- A **2-rig** is a cocomplete monoidal category where the tensor product distributes over the colimits [5], though for the purpose of constructing a compact closed bicategory we only need the tensor product to distribute over finite coproducts. Given a symmetric 2-rig R , $\text{Mat}(R)$ is the compact closed bicategory of finitely-generated free R -modules, where the tensor product is the usual tensor product for matrices. A weak monoid object in $\text{Mat}(R)$ is a categorified finite-dimensional associative algebra over R . A monad in $\text{Mat}(R)$ is a finite R -enriched category. A **finite 2-rig** S is only finitely cocomplete, but $\text{Mat}(S)$ is still compact closed. Kapranov and Voevodsky [33] described a bicategory equivalent to $\text{Mat}(\text{FinVect})$ and called its objects “2-vector spaces”.
- For a category C , let $\widehat{C} = \text{Set}^{C^{\text{op}}}$ be the category of presheaves on C . The 2-category Cocont has

– small categories as objects;

- cocontinuous functors $f : \widehat{C} \rightarrow \widehat{D}$ between the categories of presheaves on the source and target as morphisms;
- natural transformations as 2-morphisms.

We can think of cocontinuous functors as being “Set-linear transformations”, since they preserve sums. Day and Street [20] proved that Cocont is a compact closed 2-category.

- Recall that a profunctor $F : C \nrightarrow R$ is a functor $F : C^{\text{op}} \times R \rightarrow \text{Set}$; we can think of profunctors as being rather like matrices, where the set $F(c, r)$ is the “matrix element” at row r and column c . Composition of profunctors is given by taking the coend of the inner coordinates, just as matrix multiplication done by summing over the inner index. Small categories, profunctors, and natural transformations form the compact closed bicategory Prof , where the tensor product is the product in Cat . A weak monoid object in Prof is a promonoidal category [19, 29]. A symmetric monoidal monad in Prof is a Freyd category, also known as an “Arrow” in the functional programming community [1, 31].

Cattani and Winskel [16] showed that Cocont and Prof are equivalent as bicategories. Though they do not explicitly state it, the equivalence they construct is symmetric monoidal; since symmetric monoidal equivalences preserve the dual, Cocont and Prof are equivalent as compact closed bicategories.

- So far the examples have been rather algebraic in flavor, but there are topological examples, too. The category $n\text{Cob}$ is the compact closed category whose
 - objects are $(n - 1)$ -dimensional manifolds and
 - morphisms are diffeomorphism classes of collared n -dimensional cobordisms between them,

where the tensor product is disjoint union. Atiyah [2] introduced the category informally in his paper defining topological quantum field theories.

Morton [44] defined the bicategory $n\text{Cob}_2$ whose

- objects are $(n - 2)$ -dimensional manifolds,
- morphisms are collared $(n - 1)$ -dimensional cobordisms, or “manifolds with boundary”, and
- 2-morphisms are diffeomorphism classes of collared n -dimensional maps of cobordisms, or “manifolds with corners”.

The collars are necessary to preserve the smoothness when composing 1- and 2-morphisms. Schommer-Pries proved a purely algebraic characterization of 2Cob_2 , essentially proving the “Baez-Dolan cobordism hypothesis” for the $n = 2$ case [3]. We expect that $n\text{Cob}_2$ will be found to be compact closed, but we will not prove it in this paper.

- In a letter to the author, John Baez defined two interesting compact closed bicategories.

A **directed multigraph** is a finite set E of edges and a finite set V of vertices equipped with functions $s, t : E \rightarrow V$ mapping each edge to its source and target. A **resistor network** is a directed multigraph equipped with a function r assigning a resistance in $(0, \infty)$ to each edge:

$$(0, \infty) \xleftarrow{r} E \xrightleftharpoons[t]{s} V$$

There are various choices one could make for a morphism of such networks; Baez defined a morphism of resistor networks to be a pair of functions ϵ, v making this diagram commute:

$$\begin{array}{ccccc} & & E & \xrightarrow{s} & V \\ & \swarrow r & \downarrow \epsilon & & \downarrow v \\ (0, \infty) & & E' & \xrightarrow{s} & V' \\ & \nwarrow r' & & & \\ & & E' & \xrightarrow{t} & V' \end{array}$$

Resistor networks and morphisms between them form a category ResNet ; this category has finite limits and colimits.

There is a compact closed bicategory $\text{Cospan}(\text{ResNet})$ with an important compact closed subcategory Circ consisting of cospans whose feet are resistor networks with no edges. A morphism in Circ is a **circuit**, a resistor network with chosen sets of input and output vertices across which one can measure a voltage drop.

3. Previous work

Compact closed categories were first defined by Kelly [38], and later studied in depth by Kelly and Laplaza [39].

Bénabou [10] defined bicategories and showed that small categories, distributors, and natural transformations form a bicategory Dist . Distributors later became more widely known as “profunctors”, so we will call that bicategory “Prof” instead. Later, Bénabou

defined closed bicategories and showed that Prof is closed [11]. He defined V -enriched profunctors when V is a cocomplete monoidal or symmetric monoidal closed category, defined V -Prof and proved that any V -enriched functor, regarded as a V -profunctor, has a right adjoint. More applications and details are in his lecture notes [9].

Kapranov and Voevodsky [33] defined braided semistrict monoidal 2-categories, but their definition left out some necessary axioms. Baez and Neuchl [4] gave an improved definition, but it was still missing a clause; Crans [18] gave the complete definition. See Baez and Langford [6] and Shulman [48] for details.

Gordon, Power, and Street [25] defined fully weak tricategories; a monoidal bicategory is a one-object tricategory.

Another name for semistrict monoidal 2-categories is “Gray monoids”, *i.e.* monoid objects in the 2-category Gray [24]. Day and Street [20] defined compact closed Gray monoids, and appealed to the coherence theorem of Gordon, Power, and Street to extend compact closedness to arbitrary bicategories. The semistrict approach is somewhat artificial when dealing with most “naturally occurring” bicategories, since the associator for composition of 1-morphisms is rarely the identity. However, this approach does apply to Cocont.

Katis, Sabadini and Walters [34] gave a precise account of the double-entry book-keeping method *partita doppia* in terms of the compact closed bicategory $\text{Span}(\text{RGraph})$; in a later paper [35], they cite a handwritten note by McCrudden for the “swallowtail” coherence law we use in this paper.

Preller and Lambek [46] generalized compact monoidal categories in a different direction. They considered a compact monoidal category to be a one-object bicategory satisfying some criteria, and then extend that definition to multiple objects. The resulting concept of “compact bicategory” is *not* what is being studied in this paper.

McCrudden [43] gave the first fully general definitions of braided, sylleptic, and symmetric monoidal bicategories. Schommer-Pries [47] gave the correct notion of a monoidal transformation between monoidal functors between monoidal bicategories.

Carboni and Walters [15] proved that V -Prof is a cartesian bicategory. Later, they showed [16] that Prof is equivalent to Cocont as a bicategory. Together with Kelly and Wood [14], they proved that any cartesian bicategory is symmetric monoidal in the sense of McCrudden.

4. Compact closed bicategories

In this section, we lay out the definition of a compact closed bicategory. First we give the definition of a bicategory, then start adding structure to it: we introduce the tensor product and monoidal unit; then we look at the different ways to move objects around each other, giving braided, sylleptic and symmetric monoidal bicategories. Next, we define closed monoidal bicategories by introducing a right pseudoadjoint to tensoring with an object; and finally we introduce duals for objects in a bicategory.

4.1. DEFINITION. A **bicategory** $\mathcal{K}$ consists of

1. a collection of **objects**
2. for each pair of objects A, B in $\mathcal{K}$, a category $\mathcal{K}(A, B)$; the objects of $\mathcal{K}(A, B)$ are called **1-morphisms**, while the morphisms of $\mathcal{K}(A, B)$ are called **2-morphisms**.
3. for each triple of objects A, B, C in $\mathcal{K}$, a **composition functor**

$$\circ_{A,B,C}: \mathcal{K}(B, C) \times \mathcal{K}(A, B) \rightarrow \mathcal{K}(A, C).$$

We will leave off the indices and write it as an infix operator.

4. for each object A in $\mathcal{K}$, an object 1_A in $\mathcal{K}(A, A)$ called the **identity 1-morphism on A** . We will often write this simply as A .
5. for each quadruple of objects A, B, C, D , a natural isomorphism called the **associator for composition**; if (f, g, h) is an object of $\mathcal{K}(C, D) \times \mathcal{K}(B, C) \times \mathcal{K}(A, B)$, then

$$\mathring{a}_{f,g,h}: (f \circ g) \circ h \rightarrow f \circ (g \circ h).$$

6. for each pair of objects A, B in $\mathcal{K}$, natural isomorphisms called **left and right unitors for composition**. If f is an object of $\mathcal{K}(A, B)$, then

$$\begin{aligned} \mathring{l}_f: B \circ f &\xrightarrow{\sim} f \\ \mathring{r}_f: f \circ A &\xrightarrow{\sim} f \end{aligned}$$

such that $\mathring{a}, \mathring{l}$, and $\mathring{r}$ satisfy the following coherence laws:

1. for all (f, g, h, j) in $\mathcal{K}(D, E) \times \mathcal{K}(C, D) \times \mathcal{K}(B, C) \times \mathcal{K}(A, B)$, the following diagram, called the **pentagon equation**, commutes:

$$\begin{array}{ccccc}
 & & ((f \circ g) \circ h) \circ j & & \\
 & \swarrow \mathring{a}_{f,g,h \circ j} & & \searrow \mathring{a}_{f \circ g, h, j} & \\
 (f \circ (g \circ h)) \circ j & & & & (f \circ g) \circ (h \circ j) \\
 \downarrow \mathring{a}_{f, g \circ h, j} & & & & \swarrow \mathring{a}_{f, g, h \circ j} \\
 f \circ ((g \circ h) \circ j) & & & & \\
 \searrow f \circ \mathring{a}_{g, h, j} & & & \swarrow & \\
 & f \circ (g \circ (h \circ j)) & & &
 \end{array}$$

2. for all (f, g) in $\mathcal{K}(B, C) \times \mathcal{K}(A, B)$ the following diagram, called the **triangle equation**, commutes:

$$\begin{array}{ccc}
 (f \circ B) \circ g & \xrightarrow{\hat{a}} & f \circ (B \circ g) \\
 \searrow \hat{r}_{f \circ g} & & \swarrow f \circ \hat{l}_g \\
 & f \circ g &
 \end{array}$$

The associator $\hat{a}$ and unitors $\hat{r}, \hat{l}$ for composition of 1-morphisms are necessary, but when we are drawing commutative diagrams of 1-morphisms they are very hard to show; fortunately, by the coherence theorem for bicategories, any consistent choice is equivalent to any other, so we leave them out.

We refer the reader to Tom Leinster’s excellent “Basic bicategories” [42] for definitions of

- morphisms of bicategories, which we call functors,
- transformations between functors, which we call pseudonatural transformations, and
- modifications between transformations.

4.2. **DEFINITION.** An **equivalence** of objects A, B in a bicategory is a pair of morphisms $f: A \rightarrow B, g: B \rightarrow A$ such that there exists an invertible 2-morphism $e: g \circ f \xrightarrow{\sim} 1_A$ and an invertible 2-morphism $i: f \circ g \xrightarrow{\sim} 1_B$.

4.3. **DEFINITION.** An **adjoint equivalence** is one in which the morphisms f, g are adjoint.

For a given morphism f , any two choices of data (g, e, i) making f an adjoint equivalence are canonically isomorphic, so any choice is as good as any other. When f, g form an adjoint equivalence, we write $g = f^\bullet$. Any equivalence can be improved to an adjoint equivalence.

We can often take a 2-morphism and “reverse” one of its edges. Given objects A, B, C, D , morphisms $f: A \rightarrow C, g: C \rightarrow D, h: D \rightarrow B, j: A \rightarrow B$ such that h is an adjoint equivalence, and a 2-morphism

$$\begin{array}{ccccc}
 & & j & & \\
 & \searrow & & \swarrow & \\
 A & & & & B \\
 & \searrow f & \Downarrow \alpha & \swarrow h & \\
 & C & \xrightarrow{g} & D &
 \end{array}$$

we can get a new 2-morphism

$$(\hat{r}(g) \circ f)(e_h \circ g \circ f)(h^\bullet \circ \alpha): h^\bullet \circ j \Rightarrow g \circ f,$$

$$\begin{array}{c}
\begin{array}{ccccc}
& & j & & \\
& \swarrow & & \searrow & \\
A & & & & B \\
& \searrow f & \Downarrow \alpha & \swarrow h & \searrow h^\bullet \\
& C & \xrightarrow{g} & D & \xrightarrow{h^\bullet} D \\
& \swarrow g & \Downarrow \check{r}(g)^{1_D} & \swarrow & \\
& & & &
\end{array}
\end{array}
=
\begin{array}{c}
\begin{array}{ccccc}
& & j & & \\
& \swarrow & & \searrow & \\
A & & & & B \\
& \searrow f & \Downarrow \alpha_1 & \swarrow & \searrow h^\bullet \\
& C & \xrightarrow{g} & D & \\
& & & &
\end{array}
\end{array}$$

where $e_h : h^\bullet \circ h \Rightarrow 1$ is the 2-morphism from the equivalence. We denote such variations of a 2-morphism by adding numeric subscripts; the number simply records the order in which we introduce them, not any information about the particular variation.

4.4. DEFINITION. A **monoidal bicategory** $\mathcal{K}$ is a bicategory in which we can “multiply” objects. It consists of the following:

- A bicategory $\mathcal{K}$.
- A **tensor product** functor $\otimes : \mathcal{K} \times \mathcal{K} \rightarrow \mathcal{K}$. This functor involves an invertible “tensorator” 2-morphism $(f \otimes g) \circ (f' \otimes g') \Rightarrow (f \circ f') \otimes (g \circ g')$ which we elide in most of the coherence equations below. The coherence theorem for monoidal bicategories implies that any 2-morphism involving the tensorator is the same no matter how it is inserted [27, Remark 3.1.6], so like the associator for composition of 1-morphisms, we leave it out.

The **Stasheff polytopes** [50] are a series of geometric figures whose vertices enumerate the ways to parenthesize the tensor product of n objects, so the number of vertices is given by the Catalan numbers; for each polytope, we have a corresponding $(n - 2)$ -morphism of the same shape with directed edges and faces:

1. The tensor product of one object A is the one object A itself.
2. The tensor product of two objects A and B is the one object $A \otimes B$.
3. There are two ways to parenthesize the product of three objects, so we have an **associator** adjoint equivalence pseudonatural in A, B, C


$$a : (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C)$$

for moving parentheses from the left pair to the right pair, where $(A \otimes B) \otimes C$ is denotes the functor


$$\otimes \circ (\otimes \times 1) : \mathcal{K}^3 \rightarrow \mathcal{K},$$

and similarly for $A \otimes (B \otimes C)$.


4. There are five ways to parenthesize the product of four objects, so we have a **pentagonator** invertible modification π relating the two different ways of moving parentheses from being clustered at the left to being clustered at the right. (Mnemonic: Pink Pentagonator.)



5. There are fourteen ways to parenthesize the product of five objects, so we have an **associahedron** equation of modifications with fourteen vertices relating the various ways of getting from the parentheses clustered at the left to clustered at the right.



The associahedron is a cube with three of its edges bevelled. It holds in the bicategory $\mathcal{K}$, where we have used juxtaposition instead of $\otimes$ for brevity and the unmarked 2-morphisms are instances of the pseudonaturality invertible modification for the associator. (Mnemonic for the rectangular invertible modifications: GReen conGRuences.)



- Just as in any monoid there is an identity element 1 , in every monoidal bicategory there is a **monoidal unit** object I . Associated to the monoidal unit are a series of morphisms—one of each dimension—that express how to “cancel” the unit in a product. Each morphism of dimension $n > 0$ has two Stasheff polytopes of dimension $n - 1$ as “subcells”, one for parenthesizing $n + 1$ objects and the other for parenthesizing the n objects left over after cancellation. There are $n + 1$ ways to insert I into n objects, so there are $n + 1$ morphisms of dimension n .

1. There is one monoidal unit object I .
2. There are two **unitors** adjoint equivalences l and r that are pseudonatural in A . The Stasheff polytopes for two objects and for one object are both points, so the unitors are line segments joining them.

$$l: I \otimes A \rightarrow A$$

$$r: A \otimes I \rightarrow A.$$

3. There are three **2-unitors** invertible modifications λ, μ , and ρ . The Stasheff polytope for three objects is a line segment and the Stasheff polytope for two objects is a point, so these modifications are triangles. (Mnemonic: *Umber Unitor*.)

$$\begin{array}{ccc}
 (I \otimes A) \otimes B & & \\
 \downarrow l \otimes B & \Rightarrow \lambda & \nearrow a \\
 A \otimes B & & I \otimes (A \otimes B)
 \end{array}$$

$$\begin{array}{ccc}
 (A \otimes I) \otimes B & & \\
 \downarrow r \otimes B & \Rightarrow \mu & \nearrow a \\
 A \otimes B & & A \otimes (I \otimes B)
 \end{array}$$


$$\begin{array}{ccc}
 (A \otimes B) \otimes I & & \\
 \downarrow r & \Rightarrow \rho & \nearrow a \\
 A \otimes B & & A \otimes (B \otimes I)
 \end{array}$$

4. There are four equations of modifications. The Stasheff polytope for four objects is a pentagon and the Stasheff polytope for three objects is a line segment, so these equations are irregular prisms with seven vertices.


$$\begin{array}{ccc}
 ((IB)C)D & \xrightarrow{a} & (IB)(CD) \\
 \downarrow aD & & \downarrow l(CD) \\
 (I(BC))D & \xrightarrow{\uparrow \pi} & B(CD) \\
 \downarrow a & & \downarrow l \\
 I((BC)D) & \xrightarrow{Ia} & I(B(CD))
 \end{array}
 =
 \begin{array}{ccc}
 ((IB)C)D & \xrightarrow{a} & (IB)(CD) \\
 \downarrow aD & & \downarrow (IC)D \\
 (I(BC))D & \xrightarrow{\uparrow \lambda^{-1}D} & (BC)D \\
 \downarrow a & & \downarrow l \\
 I((BC)D) & \xrightarrow{Ia} & I(B(CD))
 \end{array}$$

$$\begin{array}{ccc}
 ((AI)C)D & \xrightarrow{a} & (AI)(CD) \\
 \downarrow aD & & \downarrow r(CD) \\
 (A(IC))D & \xrightarrow{\uparrow \pi} & A(ICD) \\
 \downarrow a & & \downarrow A(lD) \\
 A((IC)D) & \xrightarrow{a} & A((IC)D)
 \end{array}
 =
 \begin{array}{ccc}
 ((AI)C)D & \xrightarrow{a} & (AI)(CD) \\
 \downarrow aD & & \downarrow (rC)D \\
 (A(IC))D & \xrightarrow{\uparrow \mu^{-1}D} & (AC)D \\
 \downarrow a & & \downarrow A(lD) \\
 A((IC)D) & \xrightarrow{a} & A((IC)D)
 \end{array}$$


- $(n = 1, k = 2)$ and $(n = 2, k = 1)$: $\binom{1+2}{1} = \binom{2+1}{1} = 3$, so whenever the associator is the identity—e.g. in a braided strictly monoidal bicategory—these polytopes are triangles, invertible modifications whose edges are the directed $(1,1)$ polytope, the braiding.




When the associator is not the identity, the triangles' vertices get replaced with associators, effectively truncating them, and we are left with hexagon invertible modifications. (Mnemonic: Blue Braiding.)




- $(n = 3, k = 1)$ and $(n = 1, k = 3)$: $\binom{3+1}{1} = \binom{1+3}{1} = 4$, so in a braided strictly monoidal bicategory, these polytopes are tetrahedra whose faces are the $(2,1)$ polytope.




Again, when the associator is not the identity, the vertices get truncated, this time being replaced by pentagonators; as a side-effect, four of the six edges are also beveled.




This equation governs shuffling one object A into three objects B, C, D :




This equation governs shuffling one object D into three objects A, B, C :




- $(n = 2, k = 2)$: $\binom{2+2}{2} = 6$; in a braided strictly monoidal bicategory, this polytope is composed mostly of $(2,1)$ triangles, but there is a pair of braidings that commute, so one face is a square.





When the associator is not the identity, the six vertices get truncated and six of the edges get beveled.



This equation governs shuffling two objects A, B into two objects C, D :





4.7. DEFINITION. In a **sytleptic** monoidal bicategory, a full twist is not necessarily equal to the identity, but may be only equivalent to it; this adjoint equivalence is called a **sytlepsis**. A sytleptic monoidal bicategory $\mathcal{K}$ is a braided monoidal bicategory equipped with a sytlepsis subject to the following axioms.

- An invertible modification (Mnemonic: Salmon Sytlepsis)

- This equation governs the interaction of the sytlepsis with the $(n = 1, k = 2)$ braiding:

- This equation governs the interaction of the sytlepsis with the $(n = 2, k = 1)$ braiding:

4.8. DEFINITION. A **symmetric monoidal bicategory** is a *syllleptic monoidal bicategory* subject to the following axiom, where the unlabeled green cells are identities:

- for all objects A and B of $\mathcal{K}$, the following equation holds:

4.9. DEFINITION. Given two bicategories $\mathcal{J}, \mathcal{K}$, two functors $L : \mathcal{J} \rightarrow \mathcal{K}$ and $R : \mathcal{K} \rightarrow \mathcal{J}$ are **pseudoadjoint** if for all $A \in \mathcal{J}, B \in \mathcal{K}$ the categories $\text{Hom}_{\mathcal{K}}(LA, B)$ and $\text{Hom}_{\mathcal{J}}(A, RB)$ are adjoint equivalent pseudonaturally in A and B .

Monoidal closed bicategories satisfy the obvious weakening of the definition of monoidal closed categories:

4.10. DEFINITION. A **monoidal closed** bicategory is one in which for every object A , the functor $- \otimes A$ has a right pseudoadjoint $A \multimap -$.

Similarly, compact closed bicategories weaken the notion of duality from compact closed categories.


4.11. DEFINITION. A **compact closed** bicategory is a symmetric monoidal bicategory in which every object has a pseudoadjoint.

This means that every object A is equipped with a (weak) **dual**, an object A^* equipped with two 1-morphisms


$$i_A : I \rightarrow A \otimes A^* \quad e_A : A^* \otimes A \rightarrow I$$

and two “zig-zag” 2-isomorphisms (Mnemonic: Yellow Yanking or Xanthic Zig-zag)


$$\zeta_A : A \Rightarrow (A \otimes e_A) \circ (i_A \otimes A)$$




$$\theta_A : A^* \Rightarrow (e_A \otimes A^*) \circ (A^* \otimes i_A)$$



such that the following “swallowtail equation” holds:



We have drawn the diagrams in a strictly monoidal compact closed bicategory for clarity; when the associator is not the identity, we truncate some corners:



5. Bicategories of spans

At the start of this paper, we saw that spans of sets form a compact closed bicategory. Street [51] suggested weakening the notion of a map of spans to hold only up to 2-isomorphism, allowing to define spans in bicategories rather than mere categories; Hoffnung [28] worked out the details.

A **span** $S : X \rightarrow Y$ in a bicategory B is a pair of morphisms with the same source: $Y \xleftarrow{q} S \xrightarrow{p} X$.

A **map of spans** f is a triple (f, α, β) consisting of a morphism $f : S \rightarrow S'$ and two invertible 2-morphisms $\alpha : p \Rightarrow p'f, \beta : q \Rightarrow q'f$:

$$\begin{array}{ccccc}
 & & S & & \\
 & q \swarrow & & \searrow p & \\
 Y & & & & X \\
 & \beta \Downarrow & f & \Downarrow \alpha & \\
 & q' \swarrow & & \searrow p' & \\
 & & S' & &
 \end{array}$$

A **map of maps of spans** is a 2-morphism $\gamma : f \Rightarrow f'$ such that $\alpha' = (p'\gamma)\alpha$ and $\beta' = (q'\gamma)\beta$. Maps and maps of maps compose in the obvious ways.

Hoffnung showed that any 2-category T with finite products and weak pullbacks (also known as iso-comma objects) gives rise to a monoidal tricategory $\text{Span}(T)$ whose

- objects are objects of T ,
- morphisms are spans in T ,
- 2-morphisms are maps of spans, and
- 3-morphisms are maps of maps of spans;

The tensor product of two spans $Y \xleftarrow{q} S \xrightarrow{p} X$ and $Y' \xleftarrow{q'} S' \xrightarrow{p'} X'$ is the span

$$Y \times Y' \xleftarrow{q \times q'} S \times S' \xrightarrow{p \times p'} X \times X'.$$


Here we show that the bicategory $\text{Span}_2(T)$ whose

- objects are objects of T ,
- morphisms are spans in T , and
- 2-morphisms are 3-isomorphism classes of maps of spans.


forms a compact closed bicategory.

5.1. LEMMA. Given spans $r = B \xleftarrow{g} R \xrightarrow{f} A$ and $s = C \xleftarrow{j} S \xrightarrow{h} B$ in T and a morphism $n : R \rightarrow S$ in T such that $hn = g$, the composite span sr is isomorphic to the span $C \xleftarrow{j^n} R \xrightarrow{f} A$.


Proof. We form the composite of the two spans using the weak pullback.



By the first part of the universal property of the weak pullback, there is a unique morphism $q : R \rightarrow SR$ such that the following equation holds.



Therefore, the following maps of spans compose to form the identity:



(Note that K^{-1} is whiskered by q , so that 2-morphism is the identity.)

Let $\omega : \pi_S^R q \pi_R^S \Rightarrow \pi_S^R SR$ be the following 2-morphism:

$$\omega = \begin{array}{ccccc} & & SR & & \\ & q \nearrow & & \searrow \pi_S^R & \\ & R & \parallel & & S \\ \pi_R^S \nearrow & & & \searrow n & \\ SR & & \Downarrow K & & \\ & \searrow SR & & \nearrow \pi_S^R & \\ & & SR & & \end{array}$$

and $\rho : \pi_R^S q \pi_R^S \Rightarrow \pi_R^S$ be the following 2-morphism:

$$\rho = \begin{array}{ccccc} & & SR & & \\ & q \nearrow & & \searrow \pi_R^S & \\ & R & \parallel & & R \\ \pi_R^S \nearrow & & & \searrow R & \\ SR & & \parallel & & \\ & \searrow SR & & \nearrow \pi_R^S & \\ & & SR & & \end{array}$$

The following equation holds:

$$\begin{array}{ccccc} & & R & & \\ & \pi_R^S \nearrow & & \searrow g & \\ & SR & \Downarrow K & \parallel & B \\ q \nearrow & & & \searrow n & \\ R & & \parallel & & \\ \pi_R^S \nearrow & & & \searrow h & \\ SR & & \Downarrow K & & S \\ & \searrow SR & & \nearrow \pi_S^R & \\ & & SR & & \end{array} = \begin{array}{ccccc} & & SR & & \\ & q \nearrow & & \searrow \pi_R^S & \\ & R & \parallel & & R \\ \pi_R^S \nearrow & & & \searrow R & \\ SR & & \parallel & & \\ & \searrow SR & & \nearrow \pi_R^S & \\ & & SR & & \\ & & \Downarrow K & & \\ & & R & & \\ & & \searrow g & & B \\ & & \parallel & & \\ & & n & & \\ & & \searrow h & & S \\ & & \nearrow \pi_S^R & & \end{array}$$

(Note on the left hand side that the K in the upper right is whiskered by q , so that 2-morphism is the identity by the definition of q .) Therefore, by the second part of the universal property of the weak pullback, there exists a unique invertible 2-morphism

$\gamma : q\pi_R^S \Rightarrow SR$ in T such that $1 = (\pi_S^R \gamma)\omega$ and $1 = (\pi_R^S \gamma)\rho$. Given γ , we can show that the following composition of maps of spans is in the same equivalence class as the identity:

$$\begin{array}{ccc}
 \begin{array}{ccccc}
 & & SR & & \\
 \pi_S^R \swarrow & & \downarrow \pi_R^S & \searrow \pi_R^S & \\
 S & \xleftarrow{n} & R & \xrightarrow{R} & R \\
 \pi_S^R \swarrow & & \downarrow q & \searrow \pi_R^S & \\
 & & SR & &
 \end{array}
 & = &
 \begin{array}{ccccc}
 & & SR & & \\
 \pi_S^R \swarrow & & \downarrow \pi_R^S & \searrow \pi_R^S & \\
 S & \xleftarrow{\downarrow K^{-1}} & R & \xrightarrow{q\pi_R^S} & R \\
 \pi_S^R \swarrow & & \downarrow & \searrow \pi_R^S & \\
 & & SR & &
 \end{array}
 \end{array}$$

$$\begin{array}{ccccc}
 & & SR & & \\
 \pi_S^R \swarrow & & \downarrow \pi_R^S & \searrow \pi_R^S & \\
 S & \xleftarrow{\downarrow K^{-1}} & R & \xrightarrow{q\pi_R^S} & R \\
 \pi_S^R \swarrow & & \downarrow & \searrow \pi_R^S & \\
 & & SR & &
 \end{array}$$

The diagram below illustrates the 3-morphism γ as a green double arrow between two spans. The left span has vertices S (left), SR (top), and SR (bottom), with arrows π_S^R , π_R^S , and $\downarrow K^{-1}$. The right span has vertices S (left), R (right), and SR (bottom), with arrows π_S^R , π_R^S , and $\downarrow$. The 3-morphism γ is represented by a green double arrow between the two spans, with red dashed lines and labels SR and $\parallel$ indicating its properties.

Since both $\pi_R^S q$ and $q\pi_R^S$ are in the same equivalence class as the corresponding identities, they are inverses and we have constructed the required isomorphism. $\square$

5.2. THEOREM. *If T is a 2-category with finite products and weak pullbacks, then $\text{Span}_2(T)$ is a compact closed bicategory.*

Proof. As noted, Hoffnung [28] showed that $\text{Span}_3(T)$ is a monoidal tricategory. We refer the reader to Hoffnung's paper for the complete definition of a monoidal tricategory, but suffice it to say that it replaces the commuting polyhedra in the above definition of a monoidal bicategory with polyhedra that commute up to a specified 3-morphism, and then adds coherence law polytopes to govern them. When we mod out by 3-isomorphism classes of maps of spans, these 3-morphisms become trivial, so $\text{Span}_2(T)$ is a monoidal bicategory.

We need to show that $\text{Span}_2(T)$ has a syllepsis that satisfies the two syllepsis coherence laws and the symmetry coherence law, to show the existence of zig-zag 2-isomorphisms, and to show that the swallowtail coherence law holds.

Because products in T are products in the underlying category of T , the universal property of the product means that the braiding σ in T is its own inverse. The syllepsis in T is the identity, so the coherence laws making T into a symmetric monoidal 2-category are trivial.

The braiding b in $\text{Span}_2(T)$ comes from the braiding σ in T :

$$b = \begin{array}{ccc} & A \times B & \\ \sigma \swarrow & & \searrow 1 \\ B \times A & & A \times B. \end{array}$$

Since σ is its own inverse, this span is also the adjoint braiding. The syllepsis in $\text{Span}_2(T)$ is the identity map of spans, so the coherence laws making $\text{Span}_2(T)$ into a sylleptic and symmetric monoidal bicategory are all therefore trivial.

The dual of an object is the object itself; the dual of a span swaps the two legs of the span; and the dual of a map of spans swaps the invertible 2-morphisms. The unit i is the span $A \times A \xleftarrow{\Delta} A \xrightarrow{!} 1$, where Δ is the diagonal 1-morphism in T and $!$ is the unique 1-morphism into the terminal object in T . The counit e is the reverse of the span i .

In order to prove that the swallowtail coherence law holds, we will show that all the 2-morphisms in the swallowtail coherence diagram are **strict isomaps of spans**—that is, they are all in the same equivalence class as a map of spans of the form

$$\begin{array}{ccccc} & & S & & \\ & q \swarrow & & \searrow p & \\ Y & & & & X \\ & 1 \Downarrow f & & 1 \Downarrow & \\ & q' \swarrow & S' & \searrow p' & \end{array}$$

where f is an isomorphism. Most of the 2-morphisms in the coherence diagram are only adjoint equivalences in $\text{Span}_3(T)$, but they become isomorphisms when we move to $\text{Span}_2(T)$.

Note that because T has finite products, a span $Y \xleftarrow{q} S \xrightarrow{p} X$ is the same as a morphism $S \xrightarrow{(p,q)} X \times Y$. If (p, q) is monic, then a strict isomap from the span to itself is necessarily the identity. When (p, q) is monic, we say the span is **jointly monic**.

The swallowtail coherence law is a map from the span $i_A = A \times A \xleftarrow{\Delta} A \xrightarrow{!} 1$ to itself; since $(!, \Delta)$ is monic, if we can show that the swallowtail is a strict isomap of spans, then it will necessarily be the identity.

The following 2-morphisms are strict isomaps of spans:

1. **Unitors for composition.** The composition of the span $r = B \xleftarrow{g} R \xrightarrow{f} A$ with the identity span on B is isomorphic to r by Lemma 5.1, where we let $S = C = B$, h and j be the identity on B , and $n = g$. The derivation for the other unitor for composition is nearly identical.
2. **Associator for composition.** Hoffnung showed [28, Prop. 4.22] that the associator for composition is a strict isomap of spans.
3. **2-morphisms in pseudonatural transformations for $\mathbf{l}, \mathbf{r}, \mathbf{a}$.** The left unitor l for tensor is a pseudonatural transformation. A pseudonatural transformation has a diagram that commutes up to a 2-morphism, which in turn is subject to various coherence laws.

$$\begin{array}{ccc}
 I \otimes A & \xrightarrow{I \otimes f} & I \otimes B \\
 \downarrow l_A & \lrcorner & \downarrow l_B \\
 A & \xrightarrow{f} & B
 \end{array}$$

For brevity we write both the product of two objects and the weak pullback of a cospan in T with juxtaposition. Taking $f = B \xleftarrow{q} F \xrightarrow{p} A$, The two composite spans in the square above are

$$\begin{array}{ccccc}
 & & (1B)(1F) & & \\
 & \swarrow \pi_{1B}^{1F} & & \searrow \pi_{1F}^{1B} & \\
 & 1B & \xleftarrow{\kappa_B^{1B,1q}} & 1F & \\
 l_B \swarrow & & \xleftarrow{1q} & & \searrow 1p \\
 B & & 1B & & 1A \\
 & \swarrow 1B & & \swarrow 1q & \\
 & & 1B & &
 \end{array}$$

$$\begin{array}{ccccc}
 & & F(1A) & & \\
 & \swarrow \pi_F^{1A} & & \searrow \pi_{1A}^F & \\
 & F & \xleftarrow{\kappa_A^{p,l_A}} & 1A & \\
 q \swarrow & & \xleftarrow{1pl_F^{-1}} & & \searrow 1A \\
 B & & A & & 1A \\
 & \swarrow p & & \swarrow l_A & \\
 & & A & &
 \end{array}$$

By Lemma 5.1, the first span is isomorphic to

$$\begin{array}{ccc}
 & 1F & \\
 l_B 1q \swarrow & & \searrow 1p \\
 B & & 1A
 \end{array}$$

Because the 2-morphism in the weak pullback is invertible, symmetry and Lemma 5.1 imply that the second span is isomorphic to

$$\begin{array}{ccc}
 & F & \\
 q \swarrow & & \searrow 1p l_F^{-1} \\
 B & & 1A
 \end{array}$$


We get a strict isomap of spans

$$\begin{array}{ccccc}
 & & 1F & & \\
 & l_B 1q \swarrow & & \searrow 1p & \\
 B & & & & 1A \\
 & \parallel & l_F & \parallel & \\
 & q \swarrow & & \searrow 1p l_F^{-1} & \\
 & & F & &
 \end{array}$$


The right unitor, pentagonator, and tensorator all follow similarly; in each case, the spans have the form required by Lemma 5.1 because the right unitor, pentagonator and tensorator in T are all natural transformations rather than pseudonatural transformations.

4. **2-morphisms λ, μ, ρ , and π**


The 2-unitor λ in $\text{Span}_2(T)$ is a map between the spans $l_A B$ and $l_{AB} a_{IAB}$. The latter is the composite span



By Lemma 5.1, this span is isomorphic to




Because the 2-morphism λ in T is trivial, we get a strict isomap of spans



The 2-morphisms μ, ρ, π all follow similarly because those 2-morphisms in T are also identities.

5. 2-morphisms ζ and θ

The 2-morphism ζ is a map from the identity on A to a morphism we will call $z = r \circ (A \otimes e) \circ a \circ (i \otimes A) \circ l^\bullet$. First we break up z into two parts; by Lemma 5.1, $a \circ (i \otimes A) \circ l^\bullet$



is isomorphic to the following span:

$$\begin{array}{ccc}
 & 1A & \\
 a \circ \Delta A \circ \Delta \circ l \swarrow & & \searrow l \\
 A(AA) & & A
 \end{array}$$

Again by Lemma 5.1, when we compose this isomorphic span with the rest of z ,

$$\begin{array}{ccccccc}
 & & A1 & \xleftarrow{A!} & AA & \xleftarrow{\Delta \circ l} & 1A \\
 & \swarrow r & & \searrow A! & \searrow A\Delta & \swarrow a \circ \Delta A \circ \Delta \circ l & \searrow l \\
 A & & A1 & & A(AA) & & A
 \end{array}$$

the result is isomorphic to the following span:

$$\begin{array}{ccc}
 & 1A & \\
 r \circ A! \circ \Delta \circ l \swarrow & & \searrow l \\
 A & & A
 \end{array}$$

The 2-morphism ζ is given by the left unitor in T followed by the two isomorphisms above. The 2-morphism θ is similar.

Given that all these maps of spans are strict isomaps of jointly monic spans, the swallowtail coherence law is a strict isomap of spans between the morphism i and itself, and is therefore the identity. $\square$

6. Proofs

6.1. THEOREM. *When C is a category with finite products and pullbacks, the bicategory $\text{Span}(C)$ of objects of C , spans in C , and maps of spans is compact closed.*

Proof. When C is a category with finite products and pullbacks, $\text{Span}(C)$ is a special case of Theorem 5.2 where all the 2-morphisms in the pseudopullbacks are identities. $\square$

6.2. THEOREM. *The bicategory Rel of sets, relations, and implications, is compact closed.*

Proof. Rel is a wide sub-bicategory of $\text{Span}(\text{Set})$ —that is, it has all sets as objects, but only some of the 1-morphisms. Relations are the jointly-monic spans, the spans for which there is at most one element whose image under the legs of the span picks out any particular pair of related elements. Implications are maps of jointly-monic spans: a function between sets with at most one element is an implication. Since all of the structure morphisms in the definition of a compact closed bicategory are jointly monic and all the 2-morphisms between those structure morphisms exist, Rel is compact closed. $\square$

6.3. THEOREM. *When R is a symmetric 2-rig, the bicategory $\text{Mat}(R)$ of natural numbers, matrices with entries in $\text{Ob}(R)$, and matrices with entries in $\text{Mor}(R)$ is compact closed.*

Proof. The composite of two matrices is given by the usual matrix product, except that we use the monoidal product in R instead of multiplication and the coproduct instead of addition. The identity matrix has the monoidal unit object in R on the diagonal and the initial object off the diagonal. Since the monoidal product in R is in general only associative up to an associator, so is the product of matrices; similarly, composing with the identity matrix only gives the same answer up to the unitor for the monoidal product in R . The pentagon and triangle equations for composition hold in $\text{Mat}(R)$ because they hold for the tensor product in R .

$\text{Mat}(R)$ is symmetric monoidal under the usual tensor product for matrices, which is associative up to the associator in R ; the monoidal unit is the object 1 up to the unitor in R . Just as with the usual notion of matrices, the tensor product of two objects m, n is mn . The pentagonator and unitors are the identity in $\text{Mat}(R)$ because they are elementwise application of the associator and unitors in R , so the associahedra and 2-unitors are trivial. The symmetry involves a permutation of the rows and columns of the matrix that is its own inverse followed by the symmetry in R . Since R is a symmetric monoidal category, braiding twice in $\text{Mat}(R)$ gives the identity, which in turn implies that the syllepsis and the coherence laws governing it are trivial. Because the associahedra and unitors are trivial, the hexagon braiding modifications are identities and the coherence laws governing the braiding are trivial.

The dual in $\text{Mat}(R)$ is the identity on objects; on matrices, it is a transposition of the matrix followed by the symmetry in R . The zig-zag morphism

$$\begin{aligned} Ae \circ iA &= (\sum_{m,n} I|n\rangle\langle nmm|)(\sum_{j,k} I|kkj\rangle\langle j|) \\ &= \sum_{j,k,m,n} II|n\rangle\langle nmm|kkj\rangle\langle j| \\ &= (\sum_j II|j\rangle\langle j|) \end{aligned}$$

is the same as the identity up to the unitor in R . The swallowtail coherence law is therefore an inverse unitor, an associator, and a unitor, *i.e.* the triangle equation in R for the pair of objects (I, I) . $\square$

6.4. THEOREM. *The bicategory Cocont is compact closed.*

Proof. Day and Street [20] proved that Cocont is compact closed; here we give a more plebeian proof. Cocont is a 2-category, and every 2-category is a bicategory whose associator and unitors for composition are identities. Recall that

- the objects are small categories,
- morphisms $f : C \rightarrow D$ are cocontinuous functors $f : \widehat{C} \rightarrow \widehat{D}$, and
- 2-morphisms are natural transformations between functors.

This is a categorification of the compact closed category FinVect , where

- objects are finite sets and
- morphisms are linear transformations between the free vector spaces on the sets.

Small categories have replaced finite sets, and cocontinuity has replaced linearity.

Cocont is a monoidal bicategory:

- $A \otimes B = A \times B$, where $\times$ is the cartesian product of categories in Cat .
- I is the terminal category.
- $a = \widehat{a^\times}$, where $a^\times$ is the associator in Cat .
- $l = \widehat{l^\times}, r = \widehat{r^\times}$, where $l^\times$ and $r^\times$ are the unitors in Cat .
- π is an equality because the associator for the cartesian product in Cat satisfies the pentagon equation.
- λ, μ, ρ are equalities because the triangle equations for the left and right unitors in Cat are equations.
- the coherence diagrams—the 14-vertex associahedron and the two 9-vertex diagrams involving the unitors—are trivial because π, λ, μ, ρ are equalities rather than merely isomorphisms.

Next, we need to check the braiding. Cocont is a braided monoidal bicategory where

- $b = \widehat{b^\times}$, where $b^\times$ is the braiding for the cartesian product in Cat .
- R and S are equalities because the braiding in Cat satisfies the hexagon equations.
- the polytopes involving R and S are trivial because R and S are equalities.

Third, we need to check the syllepsis.

- Because the braiding in Cat is symmetric, Cocont 's braiding is also symmetric;
- the syllepsis v is an equality, and

- the polytopes involving v are trivial.

Finally, to show it is compact closed, we need to check the existence of duals and to verify the swallowtail coherence law.

- Every small category A has the dual $A^* = A^{\text{op}}$.
- the counit is

$$\begin{aligned} e_A : \widehat{A^{\text{op}} \times A} &\rightarrow \widehat{1} \\ F &\mapsto \int^A F \\ \alpha &\mapsto \int^A \alpha \end{aligned}$$

where $\int^A F$ is the coend of F —that is, the coequalizer of the diagram

$$\begin{aligned} &\coprod_{g:a \rightarrow a'} F(a', a) \\ &F(g, a) \downarrow \quad \downarrow F(a', g) \\ &\coprod_a F(a, a). \end{aligned}$$

- the unit is

$$\begin{aligned} i_A : \widehat{1} &\rightarrow \widehat{A \times A^{\text{op}}} \\ 1 &\mapsto \text{hom} \\ h : S \rightarrow T &\mapsto \text{hom}_h : \text{hom}_S \rightarrow \text{hom}_T \end{aligned}$$

where

$$\begin{aligned} \text{hom}_S : A^{\text{op}} \times A &\rightarrow \text{Set} \\ (a, a') &\mapsto S \times \text{hom}(a, a') \\ (f, f') &\mapsto S \times \text{hom}(f, f') \end{aligned}$$

- The 2-morphism ζ is a natural isomorphism

$$\zeta_a : a \rightarrow \int^b \text{hom}(a, b) \times b.$$

It is canonical in the sense that we do not need to make any arbitrary choices to construct it. An elegant proof that it is canonical is the following one, shown to the

author by Tom Leinster.

$$\begin{aligned}
& \text{Set}\left(\int^b \text{hom}(a, b) \times F(b), S\right) \\
& \cong \int_b \text{Set}(\text{hom}(a, b) \times F(b), S) \\
& \cong \int_b \text{Set}(\text{hom}(a, b), \text{Set}(F(b), S)) \\
& \cong \text{Nat}(\text{hom}(a, -), \text{Set}(F(-), S)) \\
& \cong \text{Set}(F(a), S)
\end{aligned}$$

The first isomorphism trades a coend for an end, which is possible because the coend was in the covariant position; the second isomorphism is “currying,” which is possible because Set is cartesian closed. The third is perhaps the most important; just as the coend quotients by a relation, the end picks out those elements that already satisfy a relation. In this case, the relation is precisely the commuting square in the definition of a natural transformation between the given functors. Finally, the last isomorphism is the ordinary Yoneda lemma. The proof for the 2-morphism θ is almost identical. $\square$

6.5. THEOREM. *The bicategories $\text{Cospan}(\text{ResNet})$ and Circ are compact closed.*

Proof. The coproduct of two resistor networks is given by juxtaposition; the pushout of a cospan $S \leftarrow R \hookrightarrow T$ of resistor networks is given by juxtaposition followed by identifying the images of R in S and T . Cospans in ResNet are spans in $\text{ResNet}^{\text{op}}$, where the coproduct and pushout become product and pullback, so $\text{Cospan}(\text{ResNet})$ is compact closed as a corollary of Theorem 6.1. Since every object is self-dual in $\text{Cospan}(\text{ResNet})$, the subcategory Circ whose objects are resistor networks with no edges is also compact closed. $\square$

Acknowledgements

Thanks to the anonymous referee for an extraordinarily careful reading; to John Baez and Mike Shulman for explanations of concepts and helpful comments; to Alex Hoffnung for his patience in explaining weak pullbacks; and to the members of the n-Category Café and MathOverflow for expository background material and references.

References

- [1] K. Asada, Arrows are strong monads, *MSFP '10 Proceedings of the third ACM SIGPLAN workshop on Mathematically structured functional programming*. (2002), 33–42. Also available at www.ipl.t.u-tokyo.ac.jp/~asada/papers/arrStrMnd.pdf.
- [2] M. Atiyah, Topological quantum field theories, *Publications Mathématiques de l'Institut des Hautes Études Scientifiques*. **68:1** (1988), 175–186.
- [3] J. Baez and J. Dolan, Higher-dimensional algebra and topological quantum field theory, *J. Math. Phys.* **36** (1995), 6073–6105. Also available at arXiv:q-alg/9503002v2.
- [4] J. Baez and M. Neuchl, Higher-dimensional algebra I: Braided monoidal 2-categories, *Adv. Math.* **121** (1996), 196–244. Also available at arXiv:q-alg/9511013.
- [5] J. Baez and J. Dolan, Higher-dimensional algebra III: n-categories and the algebra of opetopes, *Adv. Math.* **135** (1998), 145–206. Also available at arXiv:q-alg/9702014.
- [6] J. Baez and L. Langford, Higher-dimensional algebra IV: 2-Tangles, *Adv. Math.* **180** (2003), 705–764. Also available at arXiv:math.QA/9811139.
- [7] J. Baez, D. Wise and A. Crans, Exotic Statistics for Strings in 4d BF Theory, *Adv. Theor. Math. Phys.* **11** (2007), 707–749. Also available at arXiv:gr-qc/0603085.
- [8] M. A. Batanin, *The Eckman-Hilton argument and higher operads*. Available at arXiv:math/0207281.
- [9] J. Bénabou, Distributors at work. Available at <http://www.mathematik.tu-darmstadt.de/~streicher/FIBR/DiWo.pdf>.
- [10] J. Bénabou, Introduction to bicategories, in *Reports of the Midwest Category Seminar*, Springer, Berlin, (1967), 1–77.
- [11] J. Bénabou, Les distributeurs, *Sem. Math. Pure Univ. Catholique de Louvain*, **33** (1973).
- [12] F. Borceux and D. Dejean, Cauchy completion in category theory, *Cahiers Topologie Géom. Différentielle Catégoriques*, **27** (1986), 133–146.
- [13] L. Breen, On the classification of 2-gerbes and 2-stacks, *Astérisque* **225** (1994), 1–160.
- [14] A. Carboni, G. M. Kelly, R. F. C. Walters, and R. J. Wood, Cartesian bicategories II, *Theory and Applications of Categories*, **19:6** (2008) 93–124. Available at <http://www.emis.de/journals/TAC/volumes/19/6/19-06abs.html>.
- [15] A. Carboni and R. F. C. Walters, Cartesian bicategories I, *J. Pure Appl. Algebra* **49** (1987), 11–32.

- [16] G. L. Cattani and G. Winskel, Profunctors, open maps and bisimulation, *Mathematical Structures in Computer Science*, **15:3** (2005) 553–614.
- [17] E. Cheng and N. Gurski, Towards an n -category of cobordisms, *Theory and Applications of Categories* **18:10** (2007) 274–302.
- [18] S. Crans, Generalized centers of braided and sylleptic monoidal 2-categories, *Adv. Math.* **136** (1998), 183–223.
- [19] B. Day. On closed categories of functors. In Saunders Mac Lane, editor, Reports of the Midwest Category Seminar, *Lecture Notes in Mathematics*, **137** (1970) 1–38.
- [20] B.J. Day and R. Street, Monoidal bicategories and Hopf algebroids, *Adv. Math.*, **129** (1997), 99–157.
- [21] M. Fiore, T. Leinster, Objects of Categories as Complex Numbers, *Adv. Math.* **190** (2005), 264–277. Available at arXiv:math/0212377.
- [22] M. Freedman, A. Kitaev, M. Larsen and Z. Wang, Topological quantum computation. Available at arXiv:quant-ph/0101025v2.
- [23] N. Gambino and J. Kock, Polynomial functors and polynomial monads, *Math. Proc. Cambridge Phil. Soc.* **154** (2013), 153–192. Also available at <http://arxiv.org/abs/0906.4931>.
- [24] J. Gray, *Formal category theory: Adjointness for 2-Categories*, Springer Lecture Notes in Mathematics **391**, Berlin, 1974.
- [25] R. Gordon, A. J. Power, and R. Street, Coherence for tricategories, *Memoirs Amer. Math. Soc.* **117** (1995) Number 558.
- [26] G. Th. Guilbaud and P. Rosenstiehl, Analyse algébrique d’un scrutin, *Mathématiques et sciences humaines* **4** (1963), 9–33.
- [27] n. Gurski, An algebraic theory of tricategories, PhD thesis, University of Chicago, (2007). Available at <http://gauss.math.yale.edu/~mg622/tricats.pdf>
- [28] A. Hoffnung, Spans in 2-categories: A monoidal tricategory. To appear. Also available at arXiv:1112.0560
- [29] R. Houston, *Linear Logic without Units*, PhD Thesis, University of Manchester (2007). Available at arxiv:1305.2231
- [30] J. Hughes, Generalising Monads to Arrows, *Science of Computer Programming* **37** (May 2000) 67–111. Preprint available at <http://www.cse.chalmers.se/~rjmh/Papers/arrows.pdf>.

- [31] B. Jacobs, C. Heunen and I. Hasuo, Categorical semantics for arrows, *Journal of Functional Programming* **19:3-4** (2009) 403–438. The results appeared in an earlier conference paper C. Heunen, B. Jacobs, Arrows, like Monads, are Monoids, *ENTCS* **158** (2006), 219–236.
- [32] A. Joyal, R. Street and D. Verity, Traced monoidal categories, *Math. Proc. of the Cambridge Phil. Soc.* **119:3** (1996) 447–468.
- [33] M. Kapranov and V. Voevodsky, 2-categories and Zamolodchikov tetrahedra equations, *Proc. Symp. Pure Math.* **56** (1994) 177–259.
- [34] P. Katis, N. Sabadini, R.F.C. Walters, On partita doppia, *Theory and Applications of Categories* (Written in 1998, to appear). Available at arXiv:0803.2429v1.
- [35] P. Katis, R.F.C. Walters, The compact closed bicategory of left adjoints, *Math. Proc. of the Cambridge Philosophical Society* **130:1** (2001) 77–87. Also available at <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.23.4825>.
- [36] G. M. Kelly, Basic concepts of enriched category theory, *Lecture Notes in Mathematics*, **64** (1982). Republished in *Reprints in Theory and Applications of Categories*, **10** (2005), 1–136. Available at <http://www.tac.mta.ca/tac/reprints/articles/10/tr10abs.html>.
- [37] G. Kelly, Coherence theorems for lax algebras and distributive laws, *Lecture Notes in Mathematics* **420**, Springer Verlag, Berlin, 1974, pp. 281–375.
- [38] G. M. Kelly, Many variable functorial calculus I, *Coherence in Categories, Lecture Notes in Math.*, **281** (1972) 66–105.
- [39] G. M. Kelly, M. L. Laplaza, Coherence for compact closed categories, *Journal of Pure and Applied Algebra* **19** (1980) 193–213.
- [40] Stephen Lack, A 2-categories companion, in *Towards Higher Categories*, eds. J. Baez and P. May, Springer, Berlin, 2009. Also available at arXiv:math/0702535.
- [41] M. Laplaza, Coherence for distributivity, *Lecture Notes in Mathematics* **281**, Springer Verlag, Berlin, 1972, pp. 29–72.
- [42] T. Leinster, Basic bicategories. (1998) Available at arXiv:math/9810017.
- [43] P. McCrudden, Balanced coalgebroids, *Theory and Applications of Categories*, **7 6** (2000), 71–147. Available at <http://www.emis.de/journals/TAC/volumes/7/n6/7-06abs.html>.
- [44] J. Morton, Double bicategories and double cospans, *Journal of Homotopy and Related Structures*, **4:1** (2009) , 389–428. Also available at arxiv:math/0611930.

- [45] J. Morton and J. Vicary, The Categorified Heisenberg Algebra I: A Combinatorial Representation. Available at [arxiv:1207.2054](https://arxiv.org/abs/1207.2054).
- [46] A. Preller, J. Lambek, Free compact 2-categories, *Mathematical Structures in Computer Science* **17** 2 (April 2007), 309–340. Also available at [ftp://ftp.math.mcgill.ca/pub/lambek/freecomcompact.pdf](http://ftp.math.mcgill.ca/pub/lambek/freecomcompact.pdf).
- [47] C. Schommer-Pries, The Classification of two-dimensional extended topological field theories, PhD thesis, UC Berkeley (2003). Available at <http://sites.google.com/site/chrissschommerpriesmath/Home/Schommer-Pries-Thesis.pdf>.
- [48] M. Shulman, Constructing symmetric monoidal bicategories. Available at [arXiv:1004.0993](https://arxiv.org/abs/1004.0993).
- [49] M. Shulman, Re: Inevitability in Mathematics, *n-Category Café* (2010). Available at https://golem.ph.utexas.edu/category/2010/06/inevitability_in_mathematics.html#c033741
- [50] J. Stasheff, Homotopy associativity of H -spaces I, II, *Trans. Amer. Math. Soc.* **108** (1963), 275–312.
- [51] R. Street, Fibrations in bicategories, *Cahiers de topologie et géométrie différentielle catégoriques*, **21** 2 (1980), 111–160. Also available at http://www.numdam.org/item?id=CTGDC_1980__21_2_111_0
- [52] R. Street, Functorial calculus in monoidal bicategories, *Applied Categorical Structures* **11** (2003) 219–227. Also available at <http://www.maths.mq.edu.au/~street/Extraord.pdf>.
- [53] R. Street, Frobenius monads and pseudomonoids, *J. Math. Physics* **45** (10) (October 2004) 3930–3948. Also available at <http://www.maths.mq.edu.au/~street/Frob.pdf>.